

## BIOSYNTHESIS OF AUXINS BY AZOTOBACTER

Muhammad Khalid, Muhammad Arshad, Azeem Khalid and Zahir Ahmad Zahir\*

### ABSTRACT

Various *Azotobacter* cultures were isolated from the rhizosphere of maize. Ten fast growing cultures were selected (numbered as Z<sub>1</sub>, Z<sub>2</sub>, .....Z<sub>10</sub>), and evaluated for their potential to produce auxins (IAA-equivalents) in vitro both in the absence and presence of auxin precursor, L-tryptophan (L-TRP). *Azotobacter* cultures varied in their auxin producing ability and Z<sub>4</sub> was the most efficient auxin (3.1 µg IAA-equivalents mL<sup>-1</sup>) producer in the absence of L-TRP. Supplementation of the medium with L-TRP @ 10<sup>-3</sup> and 10<sup>-4</sup>M stimulated auxin production by almost 2-fold over the unamended control. Various environmental factors strongly affected L-TRP-dependent auxin biosynthesis in soil, and optimal productions were found at 6.0 mg L-TRP g<sup>-1</sup> soil, 6.0 mg glucose g<sup>-1</sup> soil, pH7.0, 40 °C temperature, shaking @ of 150 rpm, no nitrogen and 48 hours of incubation. Application of antibiotics (streptomycin or erythromycin) had inhibitory effects on auxin biosynthesis in soil.